



Bright Futures Parent Handout 2 Week Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

How You Are Feeling

- Taking care of yourself gives you the energy to care for your baby. Remember to go for your postpartum checkup.
- Call for help if you feel sad or blue, or very tired for more than a few days.
- Know that returning to work or school is hard for many parents.
- Find safe, loving child care for your baby. You can ask us for help.
- If you plan to go back to work or school, start thinking about how you can keep breastfeeding.

PARENTAL WELL-BEING

Getting to Know Your Baby

- Have simple routines each day for bathing, feeding, sleeping, and playing.
- Put your baby to sleep on his back.
 - In your room.
 - Not in your bed.
 - In a crib, with slats less than 2 3/8 inches apart.
 - With the crib's sides always up.
- If using a playpen, make sure the weave is less than 1/4 inch and never leave the baby in it with the drop side down.
- Hold and cuddle your baby often.
 - Tummy time—put your baby on his tummy when awake and you are there to watch.
- Crying is normal and may increase when your baby is 6–8 weeks old.
- When your baby is crying, comfort him by talking, patting, stroking, and rocking.
- *Never shake your baby.*
- If you feel upset, put your baby in a safe place; call for help.

INFANT ADJUSTMENT

Safety

- Use a rear-facing car safety seat in all vehicles.
- Never put your baby in the front seat of a vehicle with a passenger air bag.
- Always wear your seat belt and never drive after using alcohol or drugs.
- Keep your car and home smoke free.
- Keep hanging cords or strings away from and necklaces and bracelets off of your baby.
- Keep a hand on your baby when changing clothes or the diaper.

SAFETY

Your Baby and Family

- Plan with your partner, friends, and family to have time for yourself.
- Take time with your partner too.
- Let us know if you are having any problems and cannot make ends meet. There are resources in our community that can help you.
- Join a new parents group or call us for help to connect to others if you feel alone and lonely.
- Call for help if you are ever hit or hurt by someone and if you and your baby are not safe at home.
- Prepare for an emergency/illness.
 - Keep a first-aid kit in your home.
 - Learn infant CPR.
 - Have a list of emergency phone numbers.
 - Know how to take your baby's temperature rectally. Call us if it is 100.4°F (38.0°C) or higher.
- Wash your hands often to help your baby stay healthy.

FAMILY ADJUSTMENT

- Pat, rock, undress, or change the diaper to wake your baby to feed.
- Feed your baby when you see signs of hunger.
 - Putting hand to mouth
 - Sucking, rooting, and fussing
- End feeding when you see signs your baby is full.
 - Turning away
 - Closing the mouth
 - Relaxed arms and hands
- Breastfeed or bottle-feed 8–12 times per day.
- Burp your baby during natural feeding breaks.
- Having 5–8 wet diapers and 3–4 stools each day shows your baby is eating well.

FEEDING ROUTINES

If Breastfeeding

- Continue to take your prenatal vitamins.
- When breastfeeding is going well (usually at 4–6 weeks), you can offer your baby a bottle or pacifier.

If Formula Feeding

- Always prepare, heat, and store formula safely. If you need help, ask us.
- Feed your baby 2 oz every 2–3 hours. If your baby is still hungry, you can feed more.
- Hold your baby so you can look at each other.
- Do not prop the bottle.

What to Expect at Your Baby's 6 Week Visit

We will talk about

- Taking care of yourself and your family
- Sleep and crib safety
- Keeping your home safe for your baby
- Getting back to work or school and finding child care
- Feeding your baby

Poison Help: 1-800-222-1222

Child safety seat inspection:

1-866-SEATCHECK; seatcheck.org



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.